


H-dibaryon from Full QCD Lattice Simulations

Takashi Inoue, Nihon Univ.

for HAL QCD Collaboration

S. Aoki	Univ. Tsukuba
T. Doi	RIKEN
T. Hatsuda	RIKEN, U.Tokyo
Y. Ikeda	Tokyo Inst. Tech.
T. I.	Nihon Univ.
N. Ishii	Univ. Tsukuba
K. Murano	RIKEN
H. Nemura	Univ. Tsukuba
K. Sasaki	Univ. Tsukuba



Introduction

★ H-dibaryon: predicted compact 6-quark ($B=2$) state

- R. L. Jaffe, Phys. Rev. Lett. 38 (1977) [MIT Bag model]
- One of most famous candidates of **exotic-hadron**.
- based on the observation of no Pauli exclusion due to 1_F nature and a large attractive contribution from one-gluon-exchange.
- Does H exist in nature? $B_H > 7$ MeV is ruled out by $\Lambda\Lambda$ He. Possibility of a shallow bound state or a resonance still remains.

★ Hyperon interaction (YN, YY int.)

- are important for phys. of super-nova, neutron star and so on.
- however, are not well known due to lack of experimental data.

★ Our purpose and goal

1. We **reveal** BB int., including existence of the H-dibaryon, directly **from QCD** by using **lattice** simulation.
2. We get deeper(or intuitive) understanding of BB int.
3. People can apply the knowledge to many physics. **=GOAL**

Plan of this talk

- Introduction

- Brief background, Our purpose and goal
- Lattice QCD Simulation – brief Introduction –
- Multi-hadron system in LQCD – our approach –

- Formulas & Setup

- NBS w.f., Potential, FAQ
- Lattice, Action and Facility,
- Five ensembles, Why $SU(3)_F$ limit ?


- Results

- BB interactions, H-dibaryon
- Hyperon interactions, H in the real world

- Summary & Outlook

Lattice QCD Simulation

$$L = -\frac{1}{4} G_{\mu\nu}^a G_a^{\mu\nu} + \bar{q} \gamma^\mu \left(i \partial_\mu - g t^a A_\mu^a \right) q - m \bar{q} q$$



Vacuum expectation value

$$\begin{aligned} \langle O(\bar{q}, q, U) \rangle &= \int dU d\bar{q} dq e^{-S(\bar{q}, q, U)} O(\bar{q}, q, U) \\ &= \int dU \det D(U) e^{-S_U(U)} O(D^{-1}(U)) \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N O(D^{-1}(U_i)) \end{aligned}$$

path integral

quark propagator

$\{U_i\}$: ensemble of gauge conf. U
generated w/ probability $\det D(U) e^{-S_U(U)}$


- ★ Well defined (regularized)
- ★ Fully non-perturbative
- ★ Manifest gauge invariance
- ★ Highly predictable

Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate.
Just need to measure the NBS w.f.
Then, potential can be extracted.
 - Demand a minimal lattice volume.
No need to extrapolate to $V=\infty$.
 - Can output many observables.


Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate. Just need to measure the NBS w.f. Then, potential can be extracted.
 - Demand a minimal lattice volume. No need to extrapolate to $V=\infty$.
 - Can output many observables.




Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate. Just need to measure the NBS w.f. Then, potential can be extracted.
 - Demand a minimal lattice volume. No need to extrapolate to $V=\infty$.
 - Can output many observables.




Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate. Just need to measure the NBS w.f. Then, potential can be extracted.
 - Demand a minimal lattice volume. No need to extrapolate to $V=\infty$.
 - Can output many observables.




Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate. Just need to measure the NBS w.f. Then, potential can be extracted.
 - Demand a minimal lattice volume. No need to extrapolate to $V=\infty$.
 - Can output many observables.



Multi-hadron system in LQCD

- Conventional : use **energy eigenstate** (eigenvalue)
 - Lüscher's finite volume method for phase-shift
 - Infinite volume extrapolation to get bound state energy
- HAL : utilize a **potential $V(r) + \dots$** from the **NBS w.f.**
 - ie. an effective theory which reproduce T matrix from QCD
 - Advantages
 - No need to separate E eigenstate. Just need to measure the NBS w.f. Then, potential can be extracted.
 - Demand a minimal lattice volume. No need to extrapolate to $V=\infty$.
 - Can output many observables.



★ We can **probe the H** in this approach!

as far as H has BB component.

Formulas and Setup

Nambu-Bethe-Salpeter w.f.

- NBS wave function

$$\psi^{(a)}(\vec{r}, t) \stackrel{\text{def}}{=} \sum_{\vec{x}} \langle 0 | B_i(\vec{x} + \vec{r}, t) \overbrace{B_j(\vec{x}, t)}^{\text{the same}} | B=2, \text{a-plet} \rangle$$

QCD generated state

$$\propto \sum_{\vec{x}} G^{(a)}(\vec{x} + \vec{r}, \vec{x}, t)$$

4-point function

$$G^{(a)}(\vec{x}, \vec{y}, t - t_0) = \langle 0 | B_i(\vec{x}, t) \underbrace{B_j(\vec{y}, t)}_{\text{sink}} \underbrace{\overline{B} \overline{B}^{(a)}(t_0)}_{\text{source}} | 0 \rangle$$

- Point type octet baryon field operator at sink

$$p_\alpha(\underline{x}) = \epsilon_{c_1 c_2 c_3} (C \gamma_5)_{\beta_1 \beta_2} \delta_{\beta_3 \alpha} u(\xi_1) d(\xi_2) u(\xi_3) \quad \text{with } \xi_i = \{c_i, \beta_i, \underline{x}\}$$

$$\Lambda_\alpha(x) = -\epsilon_{c_1 c_2 c_3} (C \gamma_5)_{\beta_1 \beta_2} \delta_{\beta_3 \alpha} \sqrt{\frac{1}{6}} [d(\xi_1) s(\xi_2) u(\xi_3) + s(\xi_1) u(\xi_2) d(\xi_3) - 2u(\xi_1) d(\xi_2) s(\xi_3)]$$

- Quark wall type BB source in the flavor irreducible rep.

e.g for flavor-singlet

$$\overline{B} \overline{B}^{(1)} = -\sqrt{\frac{1}{8}} \overline{\Lambda} \overline{\Lambda} + \sqrt{\frac{3}{8}} \overline{\Sigma} \overline{\Sigma} + \sqrt{\frac{4}{8}} \overline{N} \overline{N}$$

Potential

S. Aoki, T. Hatsuda, N. Ishii, Prog. Theo. Phys. 123 89(2010)

N. Ishii et al. [HAL QCD coll.] in preparation

NBS wave function $\psi(\vec{r}, t) = \phi_{Gr}(\vec{r})e^{-E_{Gr}t} + \phi_{1st}(\vec{r})e^{-E_{1st}t} \dots$

DEFINE a “potential” through the “Schrödinger eq.” for E-eigen-sates.

$$\left[2M_B - \frac{\nabla^2}{2\mu}\right]\phi_{Gr}(\vec{r})e^{-E_{Gr}t} + \int d^3\vec{r}' U(\vec{r}, \vec{r}')\phi_{Gr}(\vec{r}')e^{-E_{Gr}t} = E_{Gr}\phi_{Gr}(\vec{r})e^{-E_{Gr}t}$$

$$\left[2M_B - \frac{\nabla^2}{2\mu}\right]\phi_{1st}(\vec{r})e^{-E_{1st}t} + \int d^3\vec{r}' U(\vec{r}, \vec{r}')\phi_{1st}(\vec{r}')e^{-E_{1st}t} = E_{1st}\phi_{1st}(\vec{r})e^{-E_{1st}t}$$

Non-local but energy independent

By adding equations

$$\left[2M_B - \frac{\nabla^2}{2\mu}\right]\psi(\vec{r}, t) + \int d^3\vec{r}' U(\vec{r}, \vec{r}')\psi(\vec{r}', t) = -\frac{\partial}{\partial t}\psi(\vec{r}, t)$$

∇ expansion & truncation

$$U(\vec{r}, \vec{r}') = \delta(\vec{r} - \vec{r}')V(\vec{r}, \nabla) = \delta(\vec{r} - \vec{r}') [V(\vec{r}) + \cancel{\nabla} + \cancel{\nabla^2} \dots]$$

Therefore, in the **leading**

$$V(\vec{r}) = \frac{1}{2\mu} \frac{\nabla^2 \psi(\vec{r}, t)}{\psi(\vec{r}, t)} - \frac{\frac{\partial}{\partial t} \psi(\vec{r}, t)}{\psi(\vec{r}, t)} - 2M_B$$

FAQ

1. Does your potential depend on the choice of **source**?
2. Does your potential depend on choice of **operator at sink**?
3. Does your potential $U(r,r')$ or $V(r)$ depend on **energy**?

FAQ

1. Does your potential depend on the choice of **source**?
 - **No**. Some sources may enhance excited states in NBS w.f. However, the contamination is not problem in our new technique.
2. Does your potential depend on choice of **operator at sink**?
 - **Yes**. It can be regarded as the “**scheme**” to define a potential. Note that a potential itself is not physical observable. We'll obtain **unique** result for observables irrespective to the choice, as long as the potential $U(r,r')$ is deduced exactly.
3. Does your potential $U(r,r')$ or $V(r)$ depend on **energy**?
 - By definition, $U(r,r')$ is non-local but energy **independent**. While, determination and validity of its leading term $V(r)$ obtained here, **depend** on energy because of the truncation. However, we know that the dependence in NN case is **very small** (thanks to our choice of sink operator = point) and **negligible** at least at $E_{\text{lab.}} = 0 - 90$ MeV. We rely on this in the following. If we find the dependence, we'll determine the next leading term form it.

Lattice, Action and Facility

β	a [fm]	Sites	L [fm]
1.83	0.121(2)	$32^3 \times 32$	3.87

- Renormalization group improved Iwasaki gauge and Non-perturbatively $O(a)$ improved Wilson quark
- We thank K.-I. [Ishikawa](#) and the [PACS-CS](#) group for providing their DDHMC/PHMC code to generate gauge configuration, and the [Columbia Physics System](#) for their lattice QCD simulation code.
- We enhance S/N of data by averaging on $4 \times 4 = 16$ source, and forward/backward propagation in time.
- All numerical computation are carried at [T2K-Tsukuba](#).



Five ensembles

T.I. et.al. Phys. Rev. Lett. 106, 162002(2011)

$SU(3)_F$ limit

$K_u = K_d = K_s$	N_cfg	M_P.S. [MeV]	M_Vec [MeV]	M_Bar [MeV]	relative to
0.13660	420	1170.9(7)	1510.4(0.9)	2274(2)	New
0.13710	360	1015.2(6)	1360.6(1.1)	2031(2)	} add cfg
0.13760	480	836.8(5)	1188.9(0.9)	1749(1)	
0.13800	360	672.3(6)	1027.6(1.0)	1484(2)	
0.13840	720	468.6(7)	829.2(1.5)	1161(2)	New

K: quark hopping parameter

- We've made five ensembles with different value of $K_u=K_d=K_s$ (=quark mass) corresponding to $M_{PS} = 1.17$ [GeV] to 470 [MeV].
- With lightest quark (bottom row of the table),
 - p.s. meson is a little lighter than the physical kaon.
 - baryon is a little lighter than the physical sigma baryon.
- Now, the simulated hadron world is not so far from the real world, although the $SU(3)_F$ breaking is not taken into account.

Why $SU(3)_F$ limit ?

- In the limit, **convenient basis** exist to describe BB int.

$$8 \times 8 = \underbrace{27 + 8s + 1}_{\text{Symmetric}} + \underbrace{10^* + 10 + 8a}_{\text{Anti-symmetric}} \quad \text{flavor irreducible rep.}$$

- In S-wave, **no off-diagonal** interaction exists.


Fermi statistics leads to

$$\begin{aligned} {}^1S_0 &: V^{(27)}(r), \quad V^{(8s)}(r), \quad V^{(1)}(r) \\ {}^3S_1 &: V^{(10^*)}(r), \quad V^{(10)}(r), \quad V^{(8a)}(r) \end{aligned}$$

- Six $V^{(a)}(r)$ contain **essential flavor-spin structure** of BB int.
- We can **reconstruct** all baryon-base interaction (eg. ΛN) by using these $V^{(a)}(r)$ with $SU(3)$ C.G. coefficients.
- The $V^{(a)}$ is useful to pin down physical origin of particular feature, since **effective models** assume the flavor symm.


Results

BB int. in flavor basis



- At the $SU(3)_F$ limit corresponding to $M_\pi = M_K = 837$ [MeV].
- QM is true at small r . Especially, **no repulsion** in $1F$ channel.
- This indicate **possibility** of a bound **H-dibaryon** in the limit.



BB int. in flavor basis



- At the $SU(3)_F$ limit corresponding to $M_\pi = M_K = 837$ [MeV].
- QM is true at small r . Especially, **no repulsion** in $1F$ channel.
- This indicate **possibility** of a bound **H-dibaryon** in the limit.

H-dibaryon



NBS w.f. & potential



- Left: Measured **NBS** w.f. of the **1F** channel
 - The finite value at large distance is **excited states** contribution as well as a finite volume effect.
- Right: Extracted **potential** of the **1F** channel
 - $V^{(1)}(r)$ become more attractive as quark mass decrease.

Observables

T. I. etal [HALQCD collaboration]
Nucl. Phys. A in print, arXive 1112.5926



- Left: scattering **phase shift** v.s. E_{cm}
 - shows existence of one discrete state below threshold.
- Right: obtained **ground state**
 - which is 20 - 50 MeV below from free BB ie. 3q-3q.
 - This means that there is a 6-quark **bound state** in the 1_F channel.
 - A stable(bound) **H-dibaryon exists** in these $SU(3)_F$ limit world! 24

Hyperon-Hyperon interaction

BB int. in baryon-basis

- In flavor SU(3) broken world, e.g. the physical one, the **baryon-basis** are used instead of the flavor-basis.
- In the SU(3)_F limit, the baryon-base potential $V_{ij}(r)$ can be obtained by a **unitary rotation** of the potential $V^{(a)}(r)$.



e.g. S=-2, l=0 sector

coupled channel

$$\begin{pmatrix} \langle \Lambda \Lambda | \\ \langle \Sigma \Sigma | \\ \langle \Xi N | \end{pmatrix} = U \begin{pmatrix} \langle 27 | \\ \langle 8 | \\ \langle 1 | \end{pmatrix}, \quad U \begin{pmatrix} V^{(27)} & & \\ & V^{(8)} & \\ & & V^{(1)} \end{pmatrix} U^t = \begin{pmatrix} V^{\Lambda\Lambda} & V^{\Lambda\Lambda}_{\Sigma\Sigma} & V^{\Lambda\Lambda}_{\Xi N} \\ & V^{\Sigma\Sigma} & V^{\Sigma\Sigma}_{\Xi N} \\ & & V^{\Xi N} \end{pmatrix}$$

- I show you potentials $V_{ij}(r)$ at the **lightest quark** mass ($K_{uds} = 0.13840$, $M_{ps}=469$ MeV) obtained with the $V^{(a)}$ in an analytic function fitted to data.

HHI in $S=-2, I=0$ sector



- $\Lambda\Lambda - N\Xi - \Sigma\Sigma$ coupled. Left: diagonal. Right: Off-diagonal.
- This sector is flavor symmetric (spin singlet), and involves $1F_7$.
- Channel coupling interactions are comparable to diagonal ones, except for small $\Lambda\Lambda - N\Xi$ transition (sign change must be artifact).
- Interaction is most attractive in $N\Xi$ channel, although it has not much meaning because channel coupling is strong.

H in the real world
– a trial calculation –


$J^P = 0^+$ states in $S=-2, I=0$

- We study scattering in $\Lambda\Lambda - N\Xi - \Sigma\Sigma$ coupled 1S_0 channel.

$$T^{\alpha\beta} = V^{\alpha\beta} + \sum_y V^{\alpha y} G_y^{(0)} T^{y\beta}, \quad G_y^{(0)} = \frac{1}{E - H_y^{(0)} + i\epsilon}$$


- For baryon masses, we use values interpolated between $SU(3)_F$ limit one at $K=0.13840$ ($M_{ps} = 469$ MeV) and physical ones linearly.

$$M_Y(x) = (1-x)M_B^{SU(3)} + xM_Y^{Phys}$$



- For $V^{\alpha\beta}$, we use LQCD results given in the previous slide.
- This is just a **trial** study or demonstration for the moment! (based on the assumptions 1. the mass of baryon has major effect, 2. qualitative features of $V^{\alpha\beta}$ remain intact w/ $SU(3)_F$ breaking),₂₉

Phase-shifts




- In the Stapp parametrization, the bar-phase-shifts are defined as

$$S_{ii}^{l=0} = \eta_i e^{2i\bar{\delta}_i}$$


- H approaches the $\Lambda\Lambda$ threshold from below and go through it.

H-dibaryon in CSM



- Energy eigenvalues in the **C**omplex-**S**caling-**M**ethod.
- H comes 3 MeV below the $N\Xi$ threshold at the empirical $SU(3)_F$ breaking in this phenomenological trial calculation.

$\Lambda\Lambda$ invariant-mass-spectrum



- Invariant-mass-spectrum of $\Lambda\Lambda$ calculated in S-wave dominance.

$$\rho_{\Lambda\Lambda}(\sqrt{s}) = |1 - S_{\Lambda\Lambda}^{l=0}|^2 / k$$

- We may have a **chance to find H** in experiments counting two Λ .³²


Summary

- ★ We've introduced motivation and purpose.
 - We want to explore for the **H-dibaryon** in **QCD** by using **Lattice**.
 - We start with **flavor SU(3) limit** to avoid complication.
- ★ We've explained our approach.
 - We extract **observables** exclusively through the interaction **potential**.
 - By using both time and spacial derivative of the **NBS w.f.** at each point, we can extract the potential even without the ground-state-saturation.
- ★ We've carried out full QCD lattice simulations for the H.
 - $32^3 \times 32$ lattice, $L=3.87$ [fm], Iwasaki gauge, clover quark
- ★ We've found a **bound=stable** H-dibaryon
 - in flavor SU(3) symmetric world at $M_{PS} = 470$ [MeV] – 1.17 [GeV],
 - its binding energy is **20 – 50** [MeV] depending on the quark mass,
- ★ We've estimated H-dibaryon in the real world.
 - w/ phenomenological SU(3)_F breaking, H may be a resonance.

Summary & Outlook

★ Plot of H binding energy from recent full QCD simulations.

- SR. Beane et al [NPLQCD colla.]
Phys. Rev. Lett. 106, 162001 (2011),
arXiv: 1109.2889[hep-lat]. ←
- Obtained binding energy from the two groups looks consistent.
- But, all data points are still away from the physical point.





- ★ We'll continue this study and put more data on this plot.
- ★ We'll take flavor SU(3) breaking into account soon.
- ★ We'll obtain information of H-dibaryon in the real world in near future.

Thank You!

Backup slides

Size of H-dibaryon



- Left: Wave function of the **lowest two** states of $V^{(1)}$.
 - the measured NBS w.f. is a **superposition** of red and green with the finite volume effect.
- Right: Comparison between the H and the physical **deuteron**.
 - One can get feeling of the H-dibaryon. **compact**.

FAQ

1. Does your potential depend on the choice of **source**?
2. Does your potential depend on choice of **operator at sink**?
3. Does your potential $U(r,r')$ or $V(r)$ depend on **energy**?

FAQ

1. Does your potential depend on the choice of **source**?
 - **No**. Some sources may enhance excited states in NBS w.f. However, the contamination is not problem in our new technique.
2. Does your potential depend on choice of **operator at sink**?
 - **Yes**. It can be regarded as the “**scheme**” to define a potential. Note that a potential itself is not physical observable. We'll obtain **unique** result for observables irrespective to the choice, as long as the potential $U(r,r')$ is deduced exactly.
3. Does your potential $U(r,r')$ or $V(r)$ depend on **energy**?
 - By definition, $U(r,r')$ is non-local but energy **independent**. While, determination and validity of its leading term $V(r)$ obtained here, **depend** on energy because of the truncation. However, we know that the dependence in NN case is **very small** (thanks to our choice of sink operator = point) and **negligible** at least at $E_{\text{lab.}} = 0 - 90$ MeV. We rely on this in the following. If we find the dependence, we'll ₃₉ determine the next leading term form it.

FAQ

4. Do you think energy dependence of $V^{(1)}(r)$ is also **small**?
5. Is the H a compact **six-quark** object or a tight **BB bound** state?

FAQ

4. Do you think energy dependence of $V^{(1)}(r)$ is also **small**?

→ **Yes**. Because a large energy dependence means that

$$\left[2M_B - \frac{\nabla^2}{2\mu} + V_{\text{Gr}}(\vec{r}) \right] \phi_{\text{Gr}}(\vec{r}) e^{-E_{\text{Gr}} t} = E_{\text{Gr}} \phi_{\text{Gr}}(\vec{r}) e^{-E_{\text{Gr}} t}$$

$$\left[2M_B - \frac{\nabla^2}{2\mu} + V_{\text{1st}}(\vec{r}) \right] \phi_{\text{1st}}(\vec{r}) e^{-E_{\text{1st}} t} = E_{\text{1st}} \phi_{\text{1st}}(\vec{r}) e^{-E_{\text{1st}} t}$$

then $V(\vec{r}) \equiv \frac{1}{2\mu} \frac{\nabla^2 \psi(\vec{r}, t)}{\psi(\vec{r}, t)} - \frac{\frac{\partial}{\partial t} \psi(\vec{r}, t)}{\psi(\vec{r}, t)} - 2M_B$ would have a large t -dep.

5. Is the H a compact **six-quark** object or a tight **BB bound** state?

→ **Both**. There is no distinct separation between two, because baryon is nothing but a 3-quark in QCD. Imagine a compact 6-quark object in $(0S)^6$ configuration. This configuration can be rewritten in a form of $(0S)^3 \times (0S)^3 \times \text{Exp}(-a r^2)$ with relative coordinate r . This shows that a compact six-quark object can have a baryonic component, which we measure in the NBS w.f. We've established existence of a stable QCD eigenstate which couples to BB state. We do NOT insist that another "H" doesn't exist which cannot couple to BB. 41

FAQ

6. Do you insist such a deeply bound H exists in the **real world**?

7. What is the **meaning** of $\sqrt{\langle r^2 \rangle}$ of H?

FAQ

6. Do you insist such a deeply bound H exists in the **real world**?
- **No**. With $SU(3)_F$ breaking, three BB thresholds in $S=-2, I=0$ sector split as $E_{\Lambda\Lambda}^{\text{Th}} < E_{N\Xi}^{\text{Th}} < E_{\Sigma\Sigma}^{\text{Th}}$. Therefore, we expect that the binding energy of H measured from $E_{\Lambda\Lambda}^{\text{Th}}$ is much smaller than the present value, or even H is above $E_{\Lambda\Lambda}^{\text{Th}}$ in the real world.
7. What is the **meaning** of $\sqrt{\langle r^2 \rangle}$ of H?
- It is a measure of spacial distribution of baryonic component in H. It corresponds to the “point matter root mean square distance” of deuteron ($2 \times 1.9 = 3.8$ [fm]).

Problem

- Free 2-body energy spectrum in **finite volume** w/ periodic B.C.

$$p_{x,y,z} = \frac{2n_{x,y,z}\pi}{L} \quad \text{then} \quad K_{n_x, n_y, n_z} = (n_x^2 + n_y^2 + n_z^2) \left(\frac{2\pi}{L} \right)^2 \frac{1}{2\mu}$$

e.g. in $L = 2$ [fm], $2\mu = M = 1750$ [MeV] case, **220** [MeV]

therefor $E_{Gr} = 2M + 0$, $E_{1st} = 2M + 220$, $E_{2nd} = 2M + 440$

- Even with interaction, spectrum is essentially the same.
- State realized in lattice at time t $|t\rangle = |Gr\rangle + |1st\rangle \dots$

$$\text{NBS w.f.} \quad \psi(\vec{r}, t) = \phi_{Gr}(\vec{r})e^{-E_{Gr}t} + \phi_{1st}(\vec{r})e^{-E_{1st}t} \dots$$

Depending on $\Delta E \equiv E_{1st} - E_{Gr}$, if t is large enough

$$\psi(\vec{r}, t) \simeq \phi_{Gr}(\vec{r})e^{-E_{Gr}t} \quad \text{Ground State Saturation} \quad \text{Exponential tail}$$

- On $L = 2$ [fm] lattice, G.S.S. is realized at $t \geq 10$ [a].

SU(3) Octet Baryon operator

$$p_\alpha = \epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} u(\xi_1) d(\xi_2) u(\xi_3)$$

$$n_\alpha = \epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} u(\xi_1) d(\xi_2) d(\xi_3)$$

$$\Sigma_\alpha^+ = -\epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} u(\xi_1) s(\xi_2) u(\xi_3)$$

$$\Sigma_\alpha^0 = -\epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} \sqrt{\frac{1}{2}} [d(\xi_1) s(\xi_2) u(\xi_3) + u(\xi_1) s(\xi_2) d(\xi_3)]$$

$$\Sigma_\alpha^- = -\epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} d(\xi_1) s(\xi_2) d(\xi_3)$$

$$\Lambda_\alpha = -\epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} \sqrt{\frac{1}{6}} [d(\xi_1) s(\xi_2) u(\xi_3) + s(\xi_1) u(\xi_2) d(\xi_3) - 2u(\xi_1) d(\xi_2) s(\xi_3)]$$

$$\Xi_\alpha^0 = \epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} s(\xi_1) u(\xi_2) s(\xi_3)$$

$$\Xi_\alpha^- = \epsilon_{c_1 c_2 c_3} (C \gamma_5)_{d_1 d_2} \delta_{d_3 \alpha} s(\xi_1) d(\xi_2) s(\xi_3)$$

- With corrected phase $\bar{1} = -\epsilon^{123} = -(ds - sd) = sd - ds$

Irreducible BB source operator

$$\overline{BB}^{(27)} = +\sqrt{\frac{27}{40}} \bar{\Lambda} \bar{\Lambda} - \sqrt{\frac{1}{40}} \bar{\Sigma} \bar{\Sigma} + \sqrt{\frac{12}{40}} \bar{N} \bar{\Xi} \quad \text{or} \quad +\sqrt{\frac{1}{2}} \bar{p} \bar{n} + \sqrt{\frac{1}{2}} \bar{n} \bar{p}$$

$$\overline{BB}^{(8s)} = -\sqrt{\frac{1}{5}} \bar{\Lambda} \bar{\Lambda} - \sqrt{\frac{3}{5}} \bar{\Sigma} \bar{\Sigma} + \sqrt{\frac{1}{5}} \bar{N} \bar{\Xi}$$

$$\overline{BB}^{(1)} = -\sqrt{\frac{1}{8}} \bar{\Lambda} \bar{\Lambda} + \sqrt{\frac{3}{8}} \bar{\Sigma} \bar{\Sigma} + \sqrt{\frac{4}{8}} \bar{N} \bar{\Xi} \quad \text{with}$$

$$\bar{\Sigma} \bar{\Sigma} = +\sqrt{\frac{1}{3}} \bar{\Sigma}^+ \bar{\Sigma}^- - \sqrt{\frac{1}{3}} \bar{\Sigma}^0 \bar{\Sigma}^0 + \sqrt{\frac{1}{3}} \bar{\Sigma}^- \bar{\Sigma}^+$$

$$\overline{BB}^{(10^*)} = +\sqrt{\frac{1}{2}} \bar{p} \bar{n} - \sqrt{\frac{1}{2}} \bar{n} \bar{p}$$

$$\bar{N} \bar{\Xi} = +\sqrt{\frac{1}{4}} \bar{p} \bar{\Xi}^- + \sqrt{\frac{1}{4}} \bar{\Xi}^- \bar{p} - \sqrt{\frac{1}{4}} \bar{n} \bar{\Xi}^0 - \sqrt{\frac{1}{4}} \bar{\Xi}^0 \bar{n}$$

$$\overline{BB}^{(10)} = +\sqrt{\frac{1}{2}} \bar{p} \bar{\Sigma}^+ - \sqrt{\frac{1}{2}} \bar{\Sigma}^+ \bar{p}$$

$$\overline{BB}^{(8a)} = +\sqrt{\frac{1}{4}} \bar{p} \bar{\Xi}^- - \sqrt{\frac{1}{4}} \bar{\Xi}^- \bar{p} - \sqrt{\frac{1}{4}} \bar{n} \bar{\Xi}^0 + \sqrt{\frac{1}{4}} \bar{\Xi}^0 \bar{n}$$

Various Theoretical Approaches to Nuclei

